

# **Cell Membrane Structure and Function**

# PREFACE

- The plasma membrane is the edge of life.
- All cells have a plasma membrane.
- It enables and maintains an internal environment of the cell that is different from its surrounding external environment.
- The plasma membrane is **selectively permeable**, it controls the traffic of molecules into and out from the cell.

**All told, each one of us has enough  
membranes in our body to cover  
about 75 soccer fields!**




# Cell Membranes

I.

Main Idea: Membranes are composed of primarily phospholipids which give the membrane a fluid like quality.

Main Idea: Membranes are embedded with proteins and sugars which give them a mosaic like quality.




Copyright © 2004 Dennis Kunkel Microscopy, Inc.

# CELLULAR MEMBRANES ARE FLUID MOSAICS OF LIPIDS AND PROTEINS

- The foundation of the membrane is the phospholipid.
- Phospholipids are **amphipathic** meaning that have a hydrophilic region and a hydrophobic region.
  - They will inherently form membranes when they come together.
- The remainder of the membrane is littered with proteins and sugars.
  - Many of the proteins are themselves amphipathic
- The **fluid mosaic model** describes the structure of membranes.




# A. Membrane Models





## B. The Fluidity of Membranes


- Membranes are static sheets, they are held together by weak hydrophobic interactions.
- The lipids and proteins both move laterally although the lipids move much more freely.





# The Fluidity of Membranes

- Membrane fluidity is directly correlated with temperature.
- The higher the temperature the more fluid the membrane
- The type of fatty acid tails in the phospholipids also effect fluidity
- Membrane fluidity is important because it effects membrane permeability




**(b) Membrane fluidity**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

# The Fluidity of Membranes

- Cholesterol acts like a membrane “fluidity buffer”.
  - At higher temperatures cholesterol decreases the membrane fluidity, by restraining lipid movement.
  - At lower temperatures cholesterol increases the membrane fluidity, by hindering the close packing.



**(c) Cholesterol within the animal cell membrane**



# C. The Evolution of Different Membrane Lipid Composition

- Variations in lipid composition appears to be an evolutionary adaptation.
- *Fish that live in cold water have a high proportion of unsaturated fatty acids.*
- *Bacteria living in thermal hot springs show a high proportion of saturated fatty acids.*





# The Evolution of Different Membrane Lipid Composition

- The ability to adjust lipid composition appears to be an evolutionary adaptation.
- *Winter wheat* can change its percentage of unsaturated fatty acids that make up its membrane as the seasons change.
- **Natural selection has favored organisms who have a mixture of fatty acid types and or those organisms that can adjust their membrane's permeability as necessary.**





# The Speed of Life

- In general, larger the beast, the slower its metabolism and the longer its life and vice versa.
- The explanation to this observation may reside in the plasma membrane.
- “Gunky” watertight membranes are found in elephants and whales with slow metabolisms and long lifespans.
  - *The most common unsaturated fatty acid accounts for 0.2% of an elephant’s membrane.*
- “Runny” leaky membranes allow mice and hummingbirds to live fast but these organisms have short lifespans.
  - *The most common unsaturated fatty acid accounts for 20% of an mouse’s membrane.*


**Can you generate hypothesis for membrane permeability and lifespan?**

# D. Membrane Proteins and Their Functions

- A membrane is a collage of different proteins.
  - More than 50 kinds of have been found so far in the plasma membrane.
- Proteins determine most of the membrane's functions .
  - Different types of cells contain different sets of membrane proteins.




# Membrane Proteins and Their Functions




**(a) Transport**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.




**(b) Enzymatic activity**




**(c) Signal transduction**


# Membrane Proteins and Their Functions



**(d) Cell-cell recognition**



**(e) Intercellular joining**



**(f) Attachment to the cytoskeleton and extracellular matrix (ECM)**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.




# The Role of Membrane Carbohydrates in Cell to Cell Recognition

- Carbohydrates play a critical role in cellular identification.
  - Very short (~15 units) sugars serve as cellular “ID tags”.
  - **Glycolipids** are sugars bound to lipids
  - **Glycoproteins** are sugars bound to proteins
- These carbs are located on the extracellular side.
  - *They vary from species to species*
  - *They vary among individuals of the same species*
  - *They vary among the cells of the same individual.*

# E. Synthesis and Sidedness of Membranes

- **Membranes have distinct inside and outside surfaces.**
- The proportion and percentage of lipids and proteins is fairly consistent even among different species
- However the types lipids and the shapes of the proteins can vary greatly




Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.



# Cell Membranes

II.

Main Idea: The structure and composition of the plasma membrane allows it to regulate the transport of substances across the membrane.




# MEMBRANE STRUCTURE RESULTS IN SELECTIVE PERMEABILITY

- Substances do not pass through the membrane indiscriminately
- The cell is able to take up some molecules while excluding others.
- Nutrients need to enter the cell and wastes need to leave the cell
- Also, substances move through the membrane at different rates



# A. The permeability of the Lipid Bilayer



# The Lipid bilayer is just one aspect of the “gatekeeper” system

**However**  
If cell permits

**Small  
Molecules**

**Fat  
Soluble  
Molecules**


**Molecules  
with no  
electrical  
charge**

**\*Water**

**Large  
Molecules**

**Water  
Soluble  
Molecules**


**Molecules  
with  
electrical  
charges**






## B. Transport Proteins

- Hydrophilic substances can pass through protein pores called ***transport proteins*** more specifically ***channel proteins*** whose channel is also hydrophilic.
- **Aquaporins** for example greatly facilitates the rate of water transport.
- Other ***transport proteins*** hold on called ***carrier proteins*** hold onto their passengers, change shape and release to the other side



(a) A channel protein



(b) A carrier protein


**Selectively permeability depends on both the discriminating nature of the lipid bilayer and the specific proteins built into the membrane.**



# Cell Membrane

## III.

Main Idea: The direction of molecular movement across a membrane is determined by concentration gradients, while the mechanism is a result of the innate movement of molecules.




# PASSIVE TRANSPORT IS DIFFUSION OF A SUBSTANCE ACROSS A MEMBRANE WITH NO ENERGY INVESTMENT

- ✱ **Passive transport** moves substances across membranes with no energy investment from the cell itself.
- ✱ However energy is still required for any movement, where does it come from?
- ✱ The *Kinetic Molecular Theory* states that molecules are in constant random motion, these molecules therefore have their own energy.
- ✱ This motion results in **diffusion**, the movement of molecules from an area of high concentration to an area of low concentration. They “spread out”!



# Diffusion




**(a) Diffusion of one solute**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

THESE YELLOW CIRCLES REPRESENT EITHER  
SOLIDS OR GASES

# Diffusion




**(b) Diffusion of two solutes**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

**DIFFUSION OF SOLUTES IS INDEPENDENT OF OTHER SOLUTES**



# A. Effects of Osmosis on Water Balance



It has been said that water moves down its concentration gradient,

However it is more accurate and helpful if focus on solute concentration and remember that water moves from a less concentrated side to a more concentrated side. OR That water moves from a low solute concentration to a high solute concentration.

# I. Water Balance of Cells Without Walls

- **Tonicity** is the ability of a surrounding solution to cause a cell to gain or lose water.
- The key is the “non-penetrating solutes” relative to cell’s interior
- **Isotonic** solutions are **same** as cell’s solution thus no net movement of water.
- **Hypertonic** solutions have **more** than a cell’s solution thus a net movement of water out of the cell.
- **Hypotonic** solutions have **less** than a cell’s solution thus a net movement of water into the cell.



## 2. Water Balance of Cells With Walls




- Because these cells have rigid walls they can not burst.
- As water moves in the cell wall pushes back (*turgor pressure*) which opposes further water uptake
- **Isotonic** solutions cause these cells to be *flaccid* (limp).
- **Hypertonic** solutions cause a net loss of water, which results in *plasmolysis* (membrane pulls away from cell wall)
  - This causes a plant to wilt and die
- **Hypotonic** solutions cause these cells to be *turgid* (stiff).
  - This is a healthy state for plants.

## Hypotonic solution




## Isotonic solution

## Hypertonic solution

**Animal cell**



**Plant cell**





cellule d'épiderme d'oignon rouge  
en présence d'eau salée.

page-svt.com

Jankowiak ced. (2009)

# Water Balance of Cells Without Walls


- As a side note:
  - Seawater is isotonic to marine invertebrates so water regulation becomes a non-issue.
  - Terrestrial animals bathe their cells in an extracellular fluid that is constantly regulated to be isotonic.
  - However many organisms must regulate their internal solutes to maintain a water balance (**osmoregulation**).
  - Recall the paramecium video from the last chapter. They use contractile vacuoles to excrete excess water.

**The water in which paramecium live is \_\_\_\_\_?**


**Hypotonic**



# B. Facilitated Diffusion



(a) A channel protein




(b) A carrier protein


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

THE ONLY DIFFERENCE BETWEEN THIS AND REGULAR DIFFUSION IS THE ROUTE...IN SIMPLE DIFFUSION MOLECULES MOVE THROUGH THE LIPID BILAYER BUT HERE MOLECULES MOVE THROUGH A CHANNEL OR CARRIER PROTEIN INSTEAD. (STILL HIGH TO LOW, STILL NO ENERGY NEEDED BY CELL)

# B. Facilitated Diffusion



(a) A channel protein



(b) A carrier protein

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.


**GATED CHANNELS:**  
**BOTH OF THESE CAN ALSO ACT LIKE GATED CHANNELS WHERE THEY OPEN OR CLOSE IN RESPONSE TO SOME STIMULI, ELECTRICAL IMPULSES FOR EXAMPLE OR THE BINDING OF A SUBSTANCE**



# Cell Membrane

## IV.

Main Idea: Molecules that do not pass freely through the lipid bilayer or protein channels can still move across the cell membrane however both an input of energy and a carrier protein is required.



# ACTIVE TRANSPORT USES ENERGY TO MOVE SOLUTES AGAINST THEIR GRADIENTS

## A. The Need for Energy in Active Transport

- **Active Transport** moves molecules against its concentration gradient and this requires the cell to invest energy.
- It also demands a *carrier protein* to carry the molecule
- **Active Transport allows the cell to maintain an internal environment different from its surroundings!**



# Internal environment that differs from the external environment




Saturday, August 24, 2013



This internal environment does not happen by accident,  
the cell must expend energy to “produce it”

**Concentration**


**Both  
Must  
Be  
Maintained**



**Charge**



# Sodium/Potassium Pump



**KEEP  
TRACK  
OF THE  
NUMBER  
OF IONS  
GOING  
IN AND  
OUT!**

# B. How Ion Pumps Maintain Membrane Potential

- ALL cells have *voltage* across their membrane, ***voltage*** is electrical potential energy
- The voltage across a membrane is called ***membrane potential***
- The cytoplasmic side is negative relative to the positive outside.
- Caused by an unequal distribution of anions and cations.
- Ranges from -50mV to -200mV (millivolts)
- Acts like a battery!





# How Ion Pumps Maintain Membrane Potential

- ***THUS two forces drive diffusion of ions across a membrane***
- A **Chemical Force** (concentration gradient)      AND
- An **Electrical Force** (membrane potential)
- The combination of forces is called the **electrochemical gradient**
- As a result we must refine our definition of diffusion
- Molecules (at least those with charges) diffuse down their electrochemical gradient.


Do you think the an electrical gradient could be strong enough to oppose the chemical gradient?

**YES**

# How Ion Pumps Maintain Membrane Potential

- Membrane proteins actively transport ions in a way that establishes a membrane potential
- The **Na<sup>+</sup>/K<sup>+</sup> pump** is the main pump used in animal cells
- The **proton pump** is the main pump used by plants, fungi and bacteria.
- **ALL cells use about 20% of their energy budget for each of these pumps thus cells use nearly half of their total energy setting up membrane potentials!**






Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

# C. Cotransport: Coupled Transport by a Membrane Protein


- A single ATP powered pump that transports a specific solute can indirectly drive the active transport of several other solutes in a mechanism called **cotransport**.






## Passive transport

### Review



## Active transport

### Review






# Cell Membrane

V.

Main Idea: Very large molecules can be moved across the membrane however it requires energy and a special mechanism.





# BULK TRANSPORT ACROSS THE PLASMA MEMBRANE OCCURS BY EXOCYTOSIS AND ENDOCYTOSIS

## A. Exocytosis



- **Exocytosis** involves the fusion of vesicles with the plasma membrane
  - Pancreas secretes insulin in this way
  - Neurons secrete acetylcholine in this way

## B. Endocytosis

- **Endocytosis** involves taking in substances by forming vesicles  
plasma membrane invaginates and pinches off
- There are three types of endocytosis:
  - *Phagocytosis* (“cell eating”)
  - *Pinocytosis* (“cell drinking”)





## PHAGOCYTOSIS



An amoeba engulfing a bacterium via phagocytosis (TEM)

## PINOCYTOSIS



Pinocytosis vesicles forming (arrows) in a cell lining a small blood vessel (TEM)