

# **A Tour of the Cell**

# PREFACE


- **Cells are fundamental to life, the cell is the simplest collection of matter than can be alive.**
- All organisms are made of cells (or a single cell)
  - Large, complex organisms are made of many cells.
  - Specialized cells working in cooperation make up these multicellular organisms.
    - Specialized cells working together form tissues, specialized tissues work together to form organs, specialized organs form systems that together carry out life functions for the complex large, organism.
- **There exists many unique cells however all cells share common features.**



# Tour of the Cell

I.

Main Idea: Biologists use microscopes and biochemistry technique called cell fractionation to study cells.



# BIOLOGISTS USE MICROSCOPES AND THE TOOLS OF BIOCHEMISTRY TO STUDY CELLS

## A. Microscopy




- Microscope Pioneers
  - 1590 Zaccharias Janssen (Dutch eyeglass maker)
  - 1665 Robert Hooke (English physicist)
  - 1670 Anton van Leeuwenhoek (Dutch, worked in dry goods store) “The Father of Microscopy”

**All the Small Things, Blink 182**



# A. Microscopy


- There 3 Important parameters in microscopy
  - **Magnification** ratio of an object's image size to its real size
  - **Resolution** clarity of the image, the minimum distance two points can be separated and still distinguished as two points
  - **Contrast** accentuates differences in parts of the sample



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

**Electron microscopes can see smaller structures than light microscopes  
HOWEVER electron microscopes can not view living organisms where as light microscopes can!**






0001


350X 1.0 kV

100u























# B. Cell Fractionation

- *Cell Fractionation* breaks cells into pieces and separates organelles and structures from each other based on their weight.
- *Centrifugation* the heaviest components will settle on the bottom of the test tube, called a pellet.
- After the “pellet” is removed faster spinning can result in smaller components collecting at the bottom (pellet)



# TECHNIQUE





**Biochemical  
Tests**

**Electron  
Microscopy**

**Reveal enzymes  
involved in cell  
respiration**

**+**

**Reveal mitochondria**

**Together this data help determine  
the function of an organelle...  
that mitochondria may carry out  
cell respiration**

# Tour of the Cell

## II.

Main Idea: There are two distinct types of cells  
*eukaryotes & prokaryotes.*

Main Idea: Prokaryotes belong to the domains of bacteria and archaea. Eukaryotes belong to protista, fungi, animals and plants.





# **EUKARYOTIC CELLS HAVE INTERNAL MEMBRANES THAT COMPARTMENTALIZE THEIR FUNCTIONS**

## **A. Comparing Eukaryotic and Prokaryotic cells**

- ALL cells share some common features:
  - plasma membranes (selective barrier)
  - cytosol (semifluid substance)
  - chromosomes (information carrier)
  - ribosomes (protein builders)

Although all cells share these common features the first and most distinct differences between cells are seen in the division between eukaryotes and prokaryotes.

- **PROKARYOTES**

- **older, less complex**
- **No nucleus (DNA in nucleoid region)**
- **No membrane bound organelles**
- **smaller**

- **EUKARYOTES**

- **younger, more complex**
- **Has nucleus (contains DNA)**
- **Has membrane bound organelles**
- **larger**

These differences are just the beginning a more comprehensive list is forth coming!



# SIZE MATTERS

The logistics to carry  
out metabolism sets the  
lower limit on cell size

The requirements for  
metabolism set the  
upper limit on cell size


**The ratio of surface area  
to volume is critical**

**Volume (cubed function) grows  
proportionately more than its surface  
area (squared function)**



**Thus a smaller object has a greater SA:V ratio**

Surface area increases while total volume remains constant




Total surface area [Sum of the surface areas (height × width) of all box sides × number of boxes]	6	150	750
Total volume [height × width × length × number of boxes]	1	125	125
Surface-to-volume (S-to-V) ratio [surface area ÷ volume]	6	1.2	6

**Bigger is Better!**



**A Big SA:V ratio is especially important for cells that continually exchange with its surroundings**

Can think of any cells or tissues that continually exchange with its environment?



**Intestine**

**Pancreas**

**Alveoli**




# I. A Panoramic View of the Eukaryotic Cell

- Eukaryotic cells have elaborately arranged internal membranes that divide the cell into compartments.
- Compartments provide different local environments that facilitate specific metabolic functions some of which are antagonistic.
- The membranes themselves are loaded with enzymes and thus participate directly in the cell's metabolism.

**Membranes are so fundamental to the organization of cells and cellular functions we will dedicate an entire chapter to them**

**COMING SOON TO A THEATER NEAR YOU**







# BioFlix


## Tour of an Animal Cell



PEARSON  
Benjamin  
Cummings

Copyright © 2007 Pearson Education, Inc., publishing as Pearson Benjamin Cummings







# BioFlix


## Tour of a Plant Cell



PEARSON  
Benjamin  
Cummings

Copyright © 2007 Pearson Education, Inc., publishing as Pearson Benjamin Cummings





The RealCaos  
Gratification Research  
[www.realcaos.com](http://www.realcaos.com)




# Tour of the Cell

## III.

Main Idea: The nucleus and ribosomes are most involved in the genetic control of a cell.

Main Idea: The nucleus contains the “blueprints” for proteins and the ribosomes serve as the “builders” of proteins.






# THE EUKARYOTIC CELL'S GENETIC INSTRUCTIONS ARE HOUSED IN THE NUCLEUS AND CARRIED OUT BY THE RIBOSOMES

## A. The Nucleus: Information Central

- The nucleus is usually the most prominent feature in a cell.
- The nucleus contains most of a cell's genes.
  - *The mitochondria and chloroplasts contain the rest.*


# The Nucleus: continued

1  $\mu\text{m}$






Surface of nuclear envelope

0.25  $\mu\text{m}$




Pore complexes (TEM)



Rough ER






**Kangaroo  
Rat  
Kidneys -  
Keratin  
filaments  
in the  
cytoplasm  
and  
nucleus of  
rat kidney  
cells help  
the cells  
maintain  
their shape  
in this**

**bovine  
pulmonary  
artery cells  
stained so  
that actin,  
mitochondria  
and DNA  
appear in  
yellow and  
blue.**



	Cell Component	Structure	Function
<b>Concept 6.3</b> The eukaryotic cell's genetic instructions are housed in the nucleus and carried out by the ribosomes		Surrounded by nuclear envelope (double membrane) perforated by nuclear pores. The nuclear envelope is continuous with the endoplasmic reticulum (ER).	Houses chromosomes, made of chromatin (DNA, the genetic material, and proteins); contains nucleoli, where ribosomal subunits are made. Pores regulate entry and exit of materials.

## The Nucleus: review

- *nuclear envelope, perforated double membrane*
- *pore complex, regulates exchange of proteins/RNA in & out*
- *nuclear lamina, protein filaments support membrane*
- *nuclear matrix, protein fibers organize genetic material*
- *chromatin, more accessible, unraveled DNA (most of time)*
- *chromosomes, highly condensed DNA packaged for replication*
- *nucleolus, synthesizes rRNA, assembles ribosomes*




## B. The Ribosomes: Protein Factories



- Complexes of rRNA and Protein
- Carrying out protein synthesis
  - Proteins are built in locations 1.) in cytoplasm or 2.) outside surface of the endoplasmic reticulum
- *Free ribosomes* assemble proteins in cytoplasm
  - these proteins are often destined to remain inside the cell
- *Bound ribosomes* assemble proteins on the surface of the endoplasmic reticulum or nuclear envelope
  - these proteins are often destined for insertion into membranes or to be exported from the cell



# Ribosomes: continued







**What trait(s) would a cell have if it possessed an above average number of ribosomes?**

**metabolically active, secretes enzymes**

**Can you think of cell types in your body that might have an above average number of ribosomes?**

**liver cells, pancreas cells, stomach cells**

**Mature red blood cells expel their nucleus, how many ribosomes would expect to find in these cells?**


**zero**



# Tour of the Cell

## IV.

Main Idea: Membrane bound organelles work together to perform a variety of important metabolic functions.





# THE ENDOMEMBRANE SYSTEM REGULATES PROTEIN TRAFFIC AND PERFORMS METABOLIC FUNCTIONS IN THE CELL

## Structures


- **Nuclear Envelope**
- **Endoplasmic Reticulum**
- **Golgi Apparatus**
- **Lysosomes**
- **Vesicles**
- **Plasma Membrane**

## Functions

- **Protein Synthesis**
- **Transportation of Proteins**
- **Chemical Reactions  
(metabolism)**
- **Transportation of Lipids**
- **Detoxification of Poisons**



# Endomembrane System







# **A. The Endoplasmic Reticulum: Biosynthetic Factory**

- **extensive network of membranes**
- **accounts for more than half of the total cell membrane**
- **two distinct types (each differs in structure and function)**
  - **Smooth E.R. (lacks ribosomes)**
  - **Rough E.R. (has ribosomes)**







# I. Functions of Smooth ER

- **Diverse functions that vary with cell type.**
- **Synthesis of lipids, *phospholipids* and *steroids***
  - *testes, ovaries, adrenal glands* produce sex hormones as a result they have an abundance of smooth E.R.
- **Enzymes that help detoxify drugs and poison**
  - adding hydroxyl groups to drugs/poison makes them more soluble and easier to secrete
  - drug addicts have more smooth E.R.
  - additional smooth E.R. contributes to their increased tolerance for drugs
- **Stores calcium in muscle cells**




## 2. Functions of Rough ER

- **The rough E.R. is still “E.R.” it has the same functions as the smooth E.R. + some!**
- The rough E.R. specializes in producing proteins that are destined for export or to become part of a membrane, they are called ***secretory proteins***
- Most *secretory proteins* are ***glycoproteins***, they have sugars attached to them
  - These proteins are kept separate from cytoplasmic proteins by the endomembrane system
- ***Transport vesicles*** move these proteins from the E.R. to their destination



# Let's take a look at this again







# B. The Golgi Apparatus: Shipping & Receiving Center

- Receives products of E.R., *these products are modified then stored or sent to their destination.*
- The Golgi stacks have distinct structural directionality
  - The E.R. side is called the *cis side... it receives*
  - The side closest to the plasma membrane is called the *trans side...it ships*



0.1  $\mu\text{m}$




TEM of Golgi apparatus



# C. Lysosomes: Digestive Compartments

- A membranous sac of hydrolytic enzymes that animal cell uses to digest macromolecules
- These hydrolytic enzymes work best in acidic environments, found in lysosomes
- **Theme: Internal environment different from external environment**

**Animal Cells Only**






Amoebas and other protists eat by engulfing smaller organisms or food particles


Nucleus

1  $\mu\text{m}$



Lysosome

White Blood Cells fight infections through phagocytosis



(a) Phagocytosis



# Cellular Recycling of macromolecules and old organelles

Human liver  
cells recycle  
half of its  
molecules  
each week


Lysosomal  
Disease:  
“Tay-Sachs”  
missing a  
lipid digesting  
enzyme in  
brain cells

Vesicle containing  
two damaged organelles

1  $\mu\text{m}$

Mitochondrion  
fragment

Peroxisome  
fragment



(b) Autophagy



# D. Vacuoles: Diverse Maintenance Compartments

- Large vesicles derived from E.R. or Golgi
  - **Theme: Internal environment different from external environment**
- Performs a variety of functions:
  - *Food vacuoles; protists (last slide)*
  - *Contractile Vacuoles; freshwater protists (use them to pump out excess water thus controlling solute concentrations)*





I Will Survive, Gloria Gaynor




# Vacuoles: continued

- Performs a variety of functions, perhaps plants and fungi show the most variation in their use of vacuoles:
  - *Hydrolytic vacuoles*; analogous to animal lysosomes
  - *Storage Vacuoles*;
    - a reserve of important organic compounds
    - a reserve of poisonous compounds for defense
    - container of pigments to attract pollinators
  - *Central Vacuoles*; a large vacuole that is repository of inorganic ions (calcium, potassium)
    - Important for plant cell growth, water is absorbed... cell enlarges with minimal investment in new cytoplasm




**Plant Cells  
Only**

Plant Cell Central Vacuole



**Can account  
for 50-70% of  
Cell Volume**




From M. I. Walker/Photo Researchers

From M. F. Brown/Visuals Unlimited



# E. The Endomembrane System: Review





# Tour of the Cell

V.

Main Idea: In Eukaryotic cells the mitochondria and chloroplasts are the organelles that convert energy into forms that the cell can use to do work.

Main Idea: The mitochondria and chloroplasts also have similar evolutionary origins.




# **MITOCHONDRIA AND CHLOROPLASTS CHANGE ENERGY FROM ONE FORM TO ANOTHER**

- ***Chloroplasts* use solar energy to build sugars from carbon dioxide and water.**
- ***Mitochondria* use the stored chemical energy in macromolecules such as sugars and fats to generate ATP (cellular fuel).**



# A. Evolutionary Origins: Mitochondria and Chloroplasts



# Origins of Mitochondria and Chloroplasts

Proposed Mitochondrial  
Ancestors



Heterotrophic Prokaryotes

Proposed Chloroplasts  
Ancestors



Autotrophic Prokaryotes

- Almost all eukaryotes have mitochondria but far less eukaryotes have chloroplasts therefore it is hypothesized that mitochondria are older than chloroplasts.
- Most speciation involves populations diverging, becoming more dissimilar to the point where they now longer can produce “fertile offspring”. A common symbol for evolution is the tree, its branches represent new lineages.
- This story is unique because two species are merging to become one new species.



# Evidence of Endosymbiosis

- Mito/Chloro are similar in size.
- Mito/Chloro similar enzymes in their membranes.
- Mito/Chloro replication resembles binary fission.
- Mito/Chloro DNA is circular.
- Mito/Chloro have their own ribosomes.
- Mito/Chloro ribosomes are the similar in size.
- Mito/Chloro rRNA sequence is similar.



**PROKARYOTES**

# B. Mitochondria: Chemical Energy Conversion


- **Found in nearly all eukaryotic cells**
  - animals, plants, fungi and most protists
- **A cell can have one, but more often has hundreds or thousands**
  - number correlates to metabolic level of the cell
- **Mitochondria are dynamic; they move, they grow and occasionally pinch into two**

Footnote: We will look at this organelle in more detail in the cell respiration unit.

**Fuel, Metallica**



We will learn more about each area and its contents and functions in the cell respiration unit



# C. Chloroplasts: Capture of Light Energy



- **Found in plants and algae**
  - chloroplasts contain a green pigment called chlorophyll
- **Chloroplasts are dynamic; they move, they grow and occasionally pinch into two**
- **Belong to the family of plastids, organelles that manufacture and store compounds. Most contain pigments. (see next slide)**

Footnote: We will look at this organelle in more detail in the cell respiration unit.

Clocks, Coldplay



# Plastids



# Comparing Mitochondria and Chloroplasts

## Mitochondria

- **Converts energy to useable forms.**
- **Not part of endomembrane system.**
- **Has double membrane.**
- **Grows and reproduces.**
- **Has its own DNA and ribosomes.**
- **Semi-autonomous.**

- 
- **Site for cell respiration**
  - **Found in both animal and plant cells**

## Chloroplasts


- **Ditto**
- **Ditto**
- **Ditto**
- **Ditto**
- **Ditto**
- **Ditto**

- 
- **Site for photosynthesis**
  - **Found in plant cells and eukaryotic algae**



# D. Peroxisomes: Oxidation

- **A Specialized Metabolic Compartment**
- **Removes hydrogen ions( $H^+$ ) from substrates and transfers them to oxygen ( $O_2$ ) thus producing hydrogen peroxide ( $H_2O_2$ )**
  - some break down fatty acids to be used in cell respiration
  - in the liver they detoxify alcohol & poisons
- Hydrogen peroxide ( $H_2O_2$ ) is itself toxic BUT there are enzymes in the peroxisomes that convert  $H_2O_2$  back to water
- **Yet another great example of the importance of compartmentalization!**



**Actin  
Microfilament  
Network**

**Stress  
Fibers**

**Nucleus**

**Peroxisomes**

**Animal Cell Micrograph**

**Figure 2**



# Tour of the Cell

## VI.

Main Idea: Protein fibers provide structure to cells and these fibers indirectly control cellular activities.




# THE CYTOSKELETON IS A NETWORK OF FIBERS THAT ORGANIZES STRUCTURES AND ACTIVITIES IN THE CELL

## A. Roles of the Cytoskeleton: Support and Motility


- Provide **support** and maintain **shape**
  - *especially important for animal cells (no cell walls)*
- **Anchors** organelles
- **Motility**; the cell itself and components within the cell








# Motility



(a)





# B. Components of the Cytoskeleton

- Thick = **Microtubules**
- Medium = **Intermediate Filaments**
- Thin = **Microfilaments**

# I. Microtubules

- Found in all eukaryotic cells



## *a. Centrosomes and Centrioles*

- *Centrosome* is a region near the nucleus, considered to be the “microtubule organizing center”
  - Not found in plants and fungi
- The *centrioles* are located within the centrosome





10  $\mu\text{m}$

Property	Microtubules (Tubulin Polymers)
Structure	Hollow tubes; wall consists of 13 columns of tubulin molecules
Diameter	25 nm with 15-nm lumen
Protein subunits	Tubulin
Main functions	Maintenance of cell shape Cell motility <div>Chromosome movements in cell division</div> <div>Organelle movements</div>











## *b. Cilia and Flagella*


- **Cellular extensions that project from the cell.**
- Cilia and Flagella have similar diameters but cilia usually occur in large numbers where flagella are limited to one or a few
- The structure of these extensions are the same
  - The microtubule arrangement is referred to as “9+2”
- Many unicellular eukaryotes use cilia or flagella for locomotion
  - Sperm of animals, plants and fungi use flagella to move
  - Cilia is often used to move fluid over a surface
  - ATP provides energy for movement



(a) Motion of flagella



5  $\mu\text{m}$




(b) Motion of cilia



15  $\mu\text{m}$






## 2. Microfilaments (Actin Filaments)


- Solid Rods shaped like a double twisted chain, also called actin
- Seems to present in all eukaryotic cells
- Known for their role in motility
  - Muscle cell contraction
  - Amoeboid movement
  - Cytoplasmic streaming



10  $\mu\text{m}$


Property	Microfilaments (Actin Filaments)
Structure	Two intertwined strands of actin
Diameter	7 nm
Protein subunits	Actin
Main functions	Maintenance of cell shape Changes in cell shape Muscle contraction Cytoplasmic streaming Cell motility Cell division







# Muscle Contraction: Sliding Filament Theory



# 3. Intermediate Filaments

- Specialized for tension bearing
- They form a more permanent framework within the cell



5  $\mu\text{m}$

**Property**

**Intermediate  
Filaments**

Structure

Fibrous proteins supercoiled  
into thicker cables

Diameter

8–12 nm

Protein  
subunits


One of several different  
proteins of the keratin family

Main  
functions

Maintenance of cell shape

Anchorage of nucleus and  
certain other organelles

Formation of nuclear lamina



**Keratin proteins**

**Fibrous subunit (keratins  
coiled together)**






# Tour of the Cell

## VII.

Main Idea: There are structures outside the cell, on its surface, that are involved in many important cellular functions.






# EXTRACELLULAR COMPONENTS AND CONNECTIONS BETWEEN CELLS HELP COORDINATE CELLULAR ACTIVITIES

## A. Cell Walls of Plants

- Cell wall lies outside plasma membrane
- It protects, shapes and prevents excess water intake.
- They are found in *plants, bacteria, fungi* and some *protists*.



**Secondary  
cell wall**

**Primary  
cell wall**

**Middle  
lamella**


1 μm



**Nature  
copied once  
more**













## Gram<sup>+</sup> Bacteria





## Gram<sup>-</sup> Bacteria



Characteristic	Gram-Positive	Gram-Negative
	 <p>LM 4 <math>\mu</math>m</p>	 <p>LM 4 <math>\mu</math>m</p>
<b>Gram Reaction</b>	Retain crystal violet dye and stain blue or purple	Can be decolorized to accept counterstain (safranin) and stain pink or red
<b>Peptidoglycan Layer</b>	Thick (multilayered)	Thin (single-layered)
<b>Teichoic Acids</b>	Present in many	Absent
<b>Periplasmic Space</b>	Absent	Present
<b>Outer Membrane</b>	Absent	Present
<b>Lipopolysaccharide (LPS) Content</b>	Virtually none	High
<b>Lipid and Lipoprotein Content</b>	Low (acid-fast bacteria have lipids linked to peptidoglycan)	High (because of presence of outer membrane)
<b>Flagellar Structure</b>	2 rings in basal body	4 rings in basal body
<b>Toxins Produced</b>	Exotoxins	Endotoxins and exotoxins
<b>Resistance to Physical Disruption</b>	High	Low
<b>Cell Wall Disruption by Lysozyme</b>	High	Low (requires pretreatment to destabilize outer membrane)
<b>Susceptibility to Penicillin and Sulfonamide</b>	High	Low
<b>Susceptibility to Streptomycin, Chloramphenicol, and Tetracycline</b>	Low	High
<b>Inhibition by Basic Dyes</b>	High	Low
<b>Susceptibility to Anionic Detergents</b>	High	Low
<b>Resistance to Sodium Azide</b>	High	Low
<b>Resistance to Drying</b>	High	Low






## **B. The Extracellular Matrix (ECM) of Animal Cells**

- Animal cells lack cell walls but they have an elaborate ECM
- The main ingredient of ECM is glycoproteins
  - The most abundant glycoprotein is collagen
  - Collagen accounts for 40% of total protein in the body
- The ECM is structurally important
- In addition the ECM can regulate cell activity, specifically it can influence the activity of genes





Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

# C. Cell Junctions


- Neighboring cells often adhere, interact, and communicate through direct physical contact.

## I. Plasmodesmata in Plant Cells

- Cell walls are perforated, cytosol can pass through adjacent cells
- These connections unify most plant cells into one living continuum.
- Specifically water, small solutes and some proteins and RNA can also pass freely through these connections



# Plasmodesma




- Apoplastic pathway (through cell wall)
- Symplastic pathway (through cytoplasm)

## 2. Tight Junctions, Desmosomes, and Gap Junctions in Animal Cells

- Three types of cell junctions
- **Tight Junctions**; proteins bind cells together very tightly
- **Desmosomes**; acts like “rivets”, fastening cells together into strong sheets
- **Gap Junctions**; provide cytoplasmic channels from one cell to another
  - proteins form pores through molecules may pass
  - essential for cell communication between certain cells like the those that make up the heart





# Comparing Animal and Plant Cells

## Animal Cells

- **Lysosomes**
- **Centrioles**
- **Flagella**
- **Gap Junctions**

## Plant Cells

- **Chloroplasts**
- **Central Vacuoles**
- **Cell wall**
- **Plasmodesmata**

---

**Most cell organelles and structures  
are shared by both cell types**



# BioVisions

at Harvard University